arXiv:0712.3359v2 [hep-ph] 9 Jan 2008

# Comment on the “extended eikonal” unitarization

S.M. Troshin

*Institute for High Energy Physics,*
*Protvino, Moscow Region, 142281, Russia*

**Abstract**

This is a comment on the recent paper by O.V. Selyugin, J.-R. Cudell, and E. Predazzi ”Analytic properties of different unitarization schemes” arXiv: 0712.0621v2, [hep-ph].

Unitarity or conservation of probability, which can be written in terms of the scattering matrix as

$$SS^{+} = 1, \tag{1}$$

implies an existence at high energies of the two scattering modes - shadowing and antishadowing. Antishadowing in the context of the rising total cross-sections and transition beyond the black disk limit was discussed in the framework of the rational unitarization scheme in [1] (Fig. 1). In the recent paper [2] we considered



S-matrix representation

$$S(s,b) = \frac{1+iU(s,b)}{1-iU(s,b)} \Longleftarrow$$

One-to-one transform

$$S \Leftrightarrow U$$

Figure 1: Rational unitarization scheme (color online)

saturation of the unitarity limit in head-on (and small impact parameter) hadron collisions, i.e. when $S(s,b)|_{b=0} \to -1$ at $s \to \infty$. It was shown that the approach to the full absorption in head-on collisions, in another words, the limit $S(s,b)|_{b=0} \to 0$ at $s \to \infty$ does not follow from unitarity, it is merely a result of the assumed saturation of the black disk limit. This limit is a direct consequence of the exponential unitarization with an extra assumption on the pure imaginary nature of the phase shift (Fig. 2). Natural interpretation of antishadowing as the



Exponential form

$S(s,b)=\exp[2i\delta(s,b)]$ $\delta(s,b)\equiv\delta_R(s,b)+i\delta_I(s,b)$

Exponential and rational forms provide $\Longrightarrow |S(s,b)|\leq 1$

$S\Leftarrow\delta$

$S\neq 0$ for any finite value of $\delta$

Figure 2: Exponential unitarization scheme (color online)

the reflective scattering was proposed. This interpretation of the unitarity saturation has grounds in the optical concepts in high energy hadron scattering. A new attempt to derive antishadowing scattering mode was made in [3]. The authors claim that “extended eikonal” unitarization scheme can also reproduce antishadowing. However, as it is evident from Fig. 3 the extended eikonal itself cannot be considered as a unitarization scheme. The authors of [3] in the new version of their paper consider possible unitarity violation when a Born amplitude has a real part and claim that such violation is small at high energies. This argument does not work at large values of impact parameters where the eikonal function is vanishingly small.



Extended eikonal form

$$\boxed{S(s,b)=2\exp[i\omega(s,b)]-1}$$

$$\xcancel{|S(s,b)|\le 1}$$

Extended eikonal form does not provide $|S(s,b)|\le 1$ and can be used (without extra constraints) only for the pure imaginary function $\omega(s,b)$

$$S\Leftarrow\omega$$

Figure 3: Extended eikonal unitarization scheme (color online)

## Acknowledgement

I am grateful to Laszlo Jenkovszky and Oleg Selyugin for the interesting and extensive correspondence.

## References

[1] S.M. Troshin, N.E. Tyurin, Phys. Lett. B **316**, 175, 1993.

[2] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. A **22**, 4437, 2007.

[3] O.V. Selyugin, J.-R. Cudell, E. Predazzi, [arXiv:0712.0621v2].